

**Stilt - Journal for the East Asian – Australasian Flyway. Number 53.  
April 2008**

IQBAL, M. 2008. **Notes on the breeding records of the White-headed Stilt in the floodplain of Ogan Komering Lebaks, South Sumatra, Indonesia.** *Stilt 53*: 3-5. (KPB-SOS, Jalang Tanjung api-api km 9 Komplek P&K Blok E1, Palembang 30152, Indonesia; EM: [kpbsos26@yahoo.com](mailto:kpbsos26@yahoo.com))

This paper provides a summary of the breeding records of White-headed Stilt *Himantopus leucocephalus* on the floodplain of the Ogan Komering Lebaks, South Sumatra, Indonesia. The floodplain is the only known breeding site of White-headed Stilt in Sumatra. The surveys confirm the presence of the White-headed Stilt on the floodplain of Ogan Komering Lebaks between June-September or during the summer and southward migration periods. Only one record was made during the winter migration period. The regular observations of nests and juveniles between 2006-2007 show that the species is a breeding visitor on the floodplain of Ogan Komering Lebaks.

ELSON, STEPHEN J. & MARCUS J.C. SINGOR. 2008. **Hooded Plover *Thinornis rubricollis tregellasi* breeding data from Western Australia.** *Stilt 53*: 6-12. (P.O. Box 25, Ongerup, WA 6336, Australia)

In Western Australia, Hooded Plover, *Thinornis rubricollis tregellasi*, breeds both on coastal beaches and at inland salt lakes. Sites investigated in this study were located at Cranbrook and Yalgorup National Park. The Cranbrook salt lake system was found to have a sedentary population of 30-40 birds and a breeding population of around six pairs. The lake acts as a refuge when the surrounding districts dry out and 330 individuals were counted in December 2007, the highest single count in Western Australia for ten years. Yalgorup National Park is located along the coast between Mandurah and Myalup and contains a number of salt lakes. It has a core breeding population of around 20 pairs and a total population that varies, depending on the season but peaks over summer with up to a maximum of 170 birds. Beach nesting sites of Hooded Plover were found from Cape Naturaliste on the Indian Ocean to Cape Arid National Park.

MELVILLE, DAVID S. & ZHANG KEJIA. 2008. **Dunlin flagged at Chongming-Dao resighted in Yukon-Kuskowin Delta, Alaska.** *Stilt 53*: 13. (Dovedale, R.D. 2 Wakefield, Nelson, New Zealand; EM: [david.melville@xtra.co.nz](mailto:david.melville@xtra.co.nz))

DUIVENVOORDEN, ANDREW. 2008. **A further instance of Hooded Plover *Thinornis rubricollis* chick predation by Nankeen Kestrel *Falco cenchroides*.** *Stilt 53*: 14. (c/- Birds Australia, Suite 2-05, 60 Leicester Street, Carlton, Vic 3053, Australia)

MINTON, CLIVE, ROSALIND JESSOP & CHRIS HASSELL. 2008. **2007 breeding success, based on juvenile ratios of Northern hemisphere Waders which spend the non-breeding season in Australia.** *Stilt 53*: 15-19. (165 Dalgetty Road, Beaumaris, Vic. 3193, Australia; EM: [mintons@ozemail.com.au](mailto:mintons@ozemail.com.au))

ROGERS, KEN G. 2008. **A note on estimating the biometrics of Bar-tailed Godwit in Australia.** *Stilt 53*: 20-23. (340 Ninks Road, St Andrews, Vic. 3761, Australia; EM: [kenrogers@hotmail.com](mailto:kenrogers@hotmail.com))

MINTON CLIVE, PENNY JOHNS & MAUREEN CHRISTIE. 2008. **Ruddy Turnstones *Arenaria interpres* on King Island, Tasmania, Australia.** *Stilt 53*: 24-29. (165 Dalgetty Road, Beaumaris, Vic. 3193, Australia; EM: [mintons@ozemail.com.au](mailto:mintons@ozemail.com.au))

WHITELAW, ANTHEA & JIM WHITELAW. 2008. **Where have all the Red-necked Stint gone?** *Stilt 53*: 30. (83 Bayview Avenue, Inverloch, Vic. 3996, Australia)

MINTON, CLIVE, ROSALIND JESSOP, CHRIS HASSELL & MAUREEN CHRISTIE. 2008. North-west Australia Wader and Tern Expedition 2007 (10<sup>th</sup> November to 1<sup>st</sup> December 2007). Stilt 53: 31-36. (165 Dalgetty Road, Beaumaris, Vic. 3193, Australia; EM: [mintons@ozemail.com.au](mailto:mintons@ozemail.com.au))